

安全 採收 期	農藥名稱	稀釋 倍數	葉 蟎 類	蚜 蟲 類	薊 馬	夜蛾 、 毒蛾 類	粉 蝨 類	斑 潛 蠅 類	切 根 蟲	施藥時期及方法
14 天	22.5%陶斯松乳劑	700							●	定植後於畦面施藥 1 次，必要時隔 10 天再施藥 1 次。
	40.8%陶斯松水基乳劑	1,000							●	
	40.8%陶斯松乳劑	1,000							●	
	22.5%陶斯松乳劑	1,500		●						害蟲發生時開始施藥，必要時隔 7 天再施藥一次。
	25%陶斯松可溼性粉劑	1,500		●						
	40.8%陶斯松乳劑	2,000		●						
	44.9%陶斯松乳劑	2,000		●						
	50%陶斯松可溼性粉劑	2,500		●						
	75%陶斯松水分散性粒劑	3,500		●						
15 天	2.5%畢芬寧水懸劑	2,000		●	●	●				
	2.8%畢芬寧乳劑	2,000	●	●	●					
	2.8%第滅寧水基乳劑	1,000		●						
	2.8%第滅寧乳劑	1,000		●						
	2.4%第滅寧水懸劑	1,000		●						
	20%達特南水溶性粒劑	2,000		●						
	3,000					●				
18 天	10%得芬瑞可溼性粉劑	3,000	●							
21 天	15%芬殺蟎水懸劑	2,000	●							
	16%可尼丁水溶性粒劑	4,000		●						
	20%芬化利水基乳劑	3,000				●				
	20%芬化利乳劑	3,000				●				
	20%芬化利可溼性粉劑	3,000				●				
	25%賽速安水溶性粒劑	5,000		●						
28 天	25%福化利乳劑	15,000		●						限花苞形成前使用
	2.5%賽洛寧微乳劑	2,000		●						
	24%福化利水基乳劑	10,000		●						
	25%福化利乳劑	10,000		●						

◎資料來源依據行政院農委會藥物毒物試驗所網站植物保護手冊，資料更新至 2014.01.09。