

抗菊病害核准登記使用藥劑一覽表

資料更新至 2014.01.09

安全採收期	農藥名稱	稀釋倍數	灰黴病	露菌病	細菌性軟腐病	疫病	莖腐病	炭疽病	白銹病	施藥時期及方法
未定	41.7%貝芬替水懸劑	1,000					●			扦插或分株後施藥一次，至全株溼潤。
	44%貝芬替水懸劑	1,000					●			
	50%貝芬替水分散性粒劑	1,200					●			
	50%貝芬替可溼性粉劑	1,200					●			
	50%貝芬替水懸劑	1,200					●			
	60%貝芬替可溼性粉劑	1,500					●			
9天	5%易胺座乳劑	1,000							●	發病初期開始施藥，必要時隔七天施藥一次。
	15%易胺座可溼性粉劑	3,000							●	
21天	40%甲基多保淨水懸劑	1,500	●							
	70%甲基多保淨可溼性粉劑	2,500	●							
	68.8%多保鏈黴素可溼性粉劑	1,000			●					
	12.5%鏈黴素溶液	1,000			●					
	23%亞托敏水懸劑	1,000				●				
		2,000		●						
	23.6%百克敏乳劑	3,000						●		
28天	40%滅派林水懸劑	4,000	●							限花苞形成前使用
	35%本達樂可溼性粉劑	2,000		●						
	50%達滅芬可溼性粉劑	3,000		●						
	18.7%達滅克敏水分散性粒劑	1,000				●				
		1,500		●						
	71.6%銅右滅達樂可溼性粉劑	600		●						
35天	62.5%賽普護汰寧水分散性粒劑	1,500	●							

◎ 資料來源依據行政院農委會藥物毒物試驗所網站植物保護手冊，資料更新至 2014.01.09。